



Rabbit anti CD146 Polyclonal Antibody

Alternative Name(s): Melanoma cell adhesion molecule (MCAM); MUC18; CD146 antigen;

Order Information

- **Description:** CD146
- **Catalogue:** 500-11134
- **Lot:** See label
- **Size:** 100ug/200ul
- **Host:** Rabbit
- **Clone:** nan
- **Application:** IHC(P), WB
- **Reactivity:** Hu

ANTIGEN PREPARATION

Recombinant human CD146 protein encoding aa 433-647 expressed in E.coli.

BACKGROUND

CD146 belongs to the immunoglobulin super-gene family with five immunoglobulin like domains (V-V-C2-C2-C2), a transmembrane region and a 63 residue cytoplasmic tail. The protein is a membrane glycoprotein that functions as a Ca²⁺ independent cell adhesion molecule involved in heterophilic cell to cell interactions. CD146 has a molecular size of 130 kDa in its reduced form (118 kDa unreduced), and N linked glycosylation accounts for fifty percent of the apparent molecular weight. In some cells the molecule carries a sulfate 3 glucuronyl moiety.

PURIFICATION

The Rabbit IgG is purified by Epitope Affinity Purification

FORMULATION

This affinity purified antibody is supplied in sterile Phosphatebuffered saline (pH7.2) containing antibody stabilizer

SPECIFICITY

This antibody reacts with human CD146. The other species are not tested.

STORAGE

The antibodies are stable for 24 months from date of receipt when stored at -20oC to -70oC. The antibodies can be stored at 2oC-8oC for three month without detectable loss of activity. Avoid repeated freezing-thawing cycles.

APPLICATIONS/SUGGESTED WORKING DILUTIONS*


- Western Blot: 0.1-1 µg/ml
- ELISA: 0.01-0.1 µg/ml
- Immunoprecipitation: 2-5 µg/ml
- IHC: 2-10 µg/ml
- Flow cytometry: Not tested
- Molecular Weight: 110.0
- Positive Control: Kidney Tissue
- Cellular Location: Cell Membrane

*Optimal dilutions should be determined by researchers for the specific applications.


FOR RESEARCH USE ONLY.

AbboMax, Inc 2528 Qume Drive, Suite 8, San Jose, California 95131, USA
1 408-573-1898 (Tel). 1 408-573-1858 (Fax). www.abbomax.com info@abbomax.com

DATA ATTACHMENTS



Western Blot: The cell lysate derived from Jurkat was immunoblotted by the Rabbit anti CD146 (Cat#500-11134) @ 1:500. An immunoreactive band around 100 kDa was observed.



Immunohistochemistry: Human Tonsil tissue (FFPE) stained with Rabbit anti-CD146 (Cat# 500-11134) at 1:50 for 30 min, RT. (Staining of formalin-fixed tissues requires boiling tissue section in 10 mM Citrate Buffer, pH 6.0 for 10 min followed by cooling at RT for 20 min.)

REFERENCES

FOR RESEARCH USE ONLY.