

Rabbit anti Estrogen Receptor (ER) (pS118) Polyclonal Antibody Alternative Name(s): Estrogen receptor 1; ESR1;

Order Information

• Description: Estrogen Receptor (ER) (pS118)

• Catalogue: 500-12264

Lot: See labelSize: 100ug/200ulHost: RabbitClone: nan

• Application: IHC(P), WB, IP

• Reactivity: Hu, Ms

ANTIGEN PREPARATION

A synthetic peptide surrounding the epitope –QL-S-PF- with a phosphorylation site of Ser118 of human estrogen receptor.

BACKGROUND

ER (estrogen receptor) belongs to a member of the steroid receptor superfamily. It contains highly conserved DNA binding (DBD) and ligand binding domains (LBD). Through its estrogen-independent and estrogen-dependent activation domains (AF-1 and AF-2, respectively), ER regulates transcription by its phosphorylation at multiple sites, such as serine 104, 106, 118, 167 and 305. Phosphorylation of certain serine at ER may relate to breast cancer process and on-going treatment

PURIFICATION

The Rabbit IgG is purified by site-modified Epitope Affinity Purification.

FORMULATION

This affinity purified antibody is supplied in sterile Tris-buffered saline (pH7.2) containing antibody stabilizer

SPECIFICITY

This antibody recognizes STAT6 (pY641) with a phosphorylated sites of Tyr641. It does not cross-react with non-phosphospecific peptide.

STORAGE

The antibodies are stable for 24 months from date of receipt when stored at -20oC to -70oC. The antibodies can be stored at 2oC-8oC for three month without detectable loss of activity. Avoid repeated freezing-thawing cycles.

APPLICATIONS/SUGGESTED WORKING DILUTIONS*

- Western Blot: 0.1-1 μg/ml
- ELISA: 0.01-0.1 μg/ml
- Immunoprecipitation: 2-5 µg/ml
- IHC: 2-10 µg/ml
- · Flow cytometry: Not tested
- Molecular Weight: 66.2
- Positive Control: Kidney Tissue
- Cellular Location: Cell Membrane

^{*}Optimal dilutions should be determined by researchers for the specific applications.



REFERENCES