



Rabbit anti c-erbB3 Polyclonal Antibody

Alternative Name(s): c-erbB3; tyrosine kinase-type cell surface receptor HER3

Order Information

- **Description:** c-erbB3
- **Catalogue:** 500-3434
- **Lot:** See label
- **Size:** 100ug/200ul
- **Host:** Rabbit
- **Clone:** nan
- **Application:** IHC(P), WB, IP
- **Reactivity:** Hu

ANTIGEN PREPARATION

A synthetic peptide corresponding to the internal segment of human c-erbB3

BACKGROUND

c-erbB3 is a member of the epidermal growth factor receptor (EGFR) family of receptor tyrosine kinases, which comprises several related receptor tyrosine kinases that are frequently overexpressed in a variety of carcinomas. Members of this receptor family include EGFR (HER1), Neu (ErbB-2, HER2), ErbB-3 (HER3) and ErbB-4 (HER4), which form either homodimers or heterodimers upon ligand binding. C-erbB3 is overexpressed in numerous cancers, including prostate, bladder, and breast tumors.

PURIFICATION

The Rabbit IgG is purified by Epitope Affinity Purification

FORMULATION

This affinity purified antibody is supplied in sterile Phosphatebuffered saline (pH7.2) containing antibody stabilizer

SPECIFICITY

This antibody recognizes human c-erbB3 . The other species not tested.

STORAGE

The antibodies are stable for 24 months from date of receipt when stored at -20°C to -70°C . The antibodies can be stored at 2°C - 8°C for three month without detectable loss of activity. Avoid repeated freezing-thawing cycles.

APPLICATIONS/SUGGESTED WORKING DILUTIONS*

- Western Blot: 0.1-1 $\mu\text{g/ml}$
- ELISA: 0.01-0.1 $\mu\text{g/ml}$
- Immunoprecipitation: 2-5 $\mu\text{g/ml}$
- IHC: 2-10 $\mu\text{g/ml}$
- Flow cytometry: Not tested
- Molecular Weight: 185.0
- Positive Control: Kidney Tissue
- Cellular Location: Cell Membrane


*Optimal dilutions should be determined by researchers for the specific applications.

FOR RESEARCH USE ONLY.

AbboMax, Inc 2528 Qume Drive, Suite 8, San Jose, California 95131, USA
1 408-573-1898 (Tel). 1 408-573-1858 (Fax). www.abbomax.com info@abbomax.com

DATA ATTACHMENTS

Western Blot: The whole lysate derived from MCF7 (20 ug/lane) immunoblotted by Rabbit anti - p53 (Cat#) at 1:500. Observed a major immunoreactive band at molecular weight ~185 kDa.



REFERENCES

FOR RESEARCH USE ONLY.