

# Rabbit anti CD34 Polyclonal Antibody

Alternative Name(s): CD34; Hematopoietic Progenitor cell antigen CD34

#### **Order Information**

- Description: CD34
- Catalogue: 500-4644
- Lot: See label
- Size: 100ug/200ul
- Host: Rabbit
- Clone: nan
- Application: IHC(P)
- Reactivity: Hu

## ANTIGEN PREPARATION

A synthetic peptide derived from the human CD34 protein.

## BACKGROUND

CD34, a single chain transmembrane glycoprotein, is selectively expressed on human lymphoid and myeloid hematopoietic progenitor cells. Staining for CD34 has been used to measure angiogenesis, which reportedly predicts tumor recurrence.

## PURIFICATION

The Rabbit IgG is purified by Epitope Affinity Purification

## FORMULATION

This affinity purified antibody is supplied in sterile Phosphatebuffered saline (pH7.2) containing antibody stabilizer

## SPECIFICITY

This antbody recognizes human CD34 . The other species not tested.

## STORAGE


The antibodies are stable for 24 months from date of receipt when stored at –20oC to –70oC. The antibodies can be stored at 2oC-8oC for three month without detectable loss of activity. Avoid repeated freezing-thawing cycles.

## **APPLICATIONS/SUGGESTED WORKING DILUTIONS\***

- Western Blot: 0.1-1 µg/ml
- ELISA: 0.01-0.1 µg/ml
- Immunoprecipitation: 2-5 µg/ml
- IHC: 2-10 µg/ml
- Flow cytometry: Not tested
- Molecular Weight: 109.0
- Positive Control: Kidney Tissue
- Cellular Location: Cell Membrane

\*Optimal dilutions should be determined by researchers for the specific applications.





Immunohistochemistry: Human Tonsil (FFPE) stained with Rabbit anti-CD34 (Cat# 500-4644) at 1:200 for 10 min @ RT. Staining of formalin-fixed tissue requires boiling tissue sections in 10 mM Citrate Buffer, pH 6.0 for 10 min followed by cooling at RT for 20 min.

## REFERENCES