



Rabbit anti JAK2(pY1007/Y1008) Polyclonal Antibody

Alternative Name(s): Janus kinase 2

Order Information

- **Description:** JAK2(pY1007/Y1008)
- **Catalogue:** 620-150
- **Lot:** See label
- **Size:** 100ug/200ul
- **Host:** Rabbit
- **Clone:** nan
- **Application:** IHC(P), WB
- **Reactivity:** Hu

ANTIGEN PREPARATION

A synthetic peptide surrounding to the epitope -EYYK- with dual phosphorylation sites at Tyr1007 and Tyr1008 of JAK2 from human, mouse and rat origins .

BACKGROUND

The Janus tyrosine kinase family is consists of four members, JAK1, JAK2, JAK3 and Tyk2. Upon the cytokine ligand binding to its the receptors, the dimerization of the receptors occurs followed by phosphorylation of the receptors. The phosphorylation of receptors recruits the tyrosine kinase physically associated to the dimerized receptors, and phosphorylate the kinase, thus activates the signal transducer and activator of thranscription (STAT1, 3, 5) and the other signal proteins.

PURIFICATION

The Rabbit IgG is purified by site-modified Epitope Affinity Purification.

FORMULATION

This affinity purified antibody is supplied in sterile Tris-buffered saline (pH7.2) containing antibody stabilizer

SPECIFICITY

This antibody recognizes JAK2 with dual phosphorylated sites of Tyr1007/1008. It does not cross-react with non-phosphospecific peptide.

STORAGE

The antibodies are stable for 24 months from date of receipt when stored at -20oC to -70oC. The antibodies can be stored at 2oC-8oC for three month without detectable loss of activity. Avoid repeated freezing-thawing cycles.

APPLICATIONS/SUGGESTED WORKING DILUTIONS*


- Western Blot: 0.1-1 µg/ml
- ELISA: 0.01-0.1 µg/ml
- Immunoprecipitation: 2-5 µg/ml
- IHC: 2-10 µg/ml
- Flow cytometry: Not tested
- Molecular Weight: 120.0
- Positive Control: Kidney Tissue
- Cellular Location: Cell Membrane

*Optimal dilutions should be determined by researchers for the specific applications.

FOR RESEARCH USE ONLY.

AbboMax, Inc 2528 Qume Drive, Suite 8, San Jose, California 95131, USA
1 408-573-1898 (Tel). 1 408-573-1858 (Fax). www.abbomax.com info@abbomax.com

DATA ATTACHMENTS



Western Blot: The tissue lysate derived from GH stimulated mouse liver was immunoblotted by Rabbit anti JAK2 (pY1007/pY1008) antibody (Cat#620-150) at 1:500. Observed a major immunoreactive band at molecular weight ~120 kDa.

REFERENCES

FOR RESEARCH USE ONLY.