

Rabbit anti LRRK2(pS1124) Polyclonal Antibody

Alternative Name(s): leucine-rich repeat kinase 2

Order Information

- Description: LRRK2(pS1124)
- Catalogue: 620-950
- Lot: See label
- Size: 100ug/200ul
- Host: Rabbit
- Clone: nan
- Application: IHC(P), WB
- Reactivity: Hu, Ms, Rt,

ANTIGEN PREPARATION

leucine S1124)

BACKGROUND

LRRK2, leucine rich repeat kinase 2, belong to the leucine-rich repeat kinase family and encodes a protein with an ankryin repeat region, a leucine-rich repeat (LRR) domain, a kinase domain, a DFG-like motif, a RAS domain, a GTPase domain, a MLK-like domain, and a WD40 domain. The protein is present largely in the cytoplasm but also associates with the mitochondrial outer membrane. Mutations in this gene have been associated with Parkinson disease-8.

PURIFICATION

The Rabbit IgG is purified by site-modified Epitope Affinity Purification.

FORMULATION

This affinity purified antibody is supplied in sterile Tris-buffered saline (pH7.2) containing antibody stabilizer

SPECIFICITY

This antibody recognizes human LRRK2(pS1124) protein with a phosphorylation site Serine 1124.

STORAGE

The antibodies are stable for 24 months from date of receipt when stored at -20oC to -70oC. The antibodies can be stored at 2oC-8oC for three month without detectable loss of activity. Avoid repeated freezing-thawing cycles.

APPLICATIONS/SUGGESTED WORKING DILUTIONS*


- Western Blot: 0.1-1 µg/ml
- ELISA: 0.01-0.1 µg/ml
- Immunoprecipitation: 2-5 µg/ml
- IHC: 2-10 µg/ml
- Flow cytometry: Not tested
- Molecular Weight: 250.0
- Positive Control: Kidney Tissue
- Cellular Location: Cell Membrane

*Optimal dilutions should be determined by researchers for the specific applications.

FOR RESEARCH USE ONLY.

AbboMax, Inc 2528 Qume Drive, Suite 8, San Jose, California 95131, USA 1 408-573-1898 (Tel). 1 408-573-1858 (Fax). www.abbomax.com info@abbomax.com





Western Blot: The whole lysate derived from PMA stimulated U118MG (20 ug/lane) immunoblotted by Rabbit anti – LRRK2 (pS1124) (Cat# 620-950) at 1:500. Observed a major immunoreactive band at molecular weight ~250kDa.

REFERENCES